



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

REC'D 09 JUN 2004

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03076044.1

BEST AVAILABLE COPY

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

PCT/EP2004 / 003995

Anmeldung Nr:
Application no.: 03076044.1
Demande no:

Anmeldetag:
Date of filing: 09.04.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Bayer BioScience N.V.
Jozef Plateaustraat 22
9000 Gent
BELGIQUE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Methods and means for increasing the tolerance of plants to stress conditions

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C12N15/82

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

Methods and means for increasing the tolerance of plants to stress conditions.

Field of the invention.

[1] The present invention relates to the use of poly (ADP-ribose) glycohydrolases in plants to increase the tolerance of plants to adverse growing conditions, including drought, high light intensities, high temperatures, nutrient limitations and the like. Methods and means are provided to produce plants that are tolerant to abiotic stress conditions.

Background to the invention

[2] . Frequently, abiotic stress will lead either directly or indirectly to damage of the DNA of the cells of the plants exposed to the adverse conditions. Genomic damage, if left unrepaired, can lead to cell death. Tolerance to stress conditions exhibited by plants is the result of the ability of the plant cells exposed to the adverse conditions to reduce and/or repair the damage, and to survive.

[3] Plant cells, like other eukaryotic cells, have evolved an elaborate DNA repair system. The activation of poly(ADP-ribose) polymerase (PARP) by DNA strand breaks is often one of the first cellular responses to DNA damage. PARP catalyzes the post-translational modification of proteins by adding successively molecules of ADP-ribose, obtained from the conversion of nicotineamide dinucleotide (NAD), to form multibranched polymers containing up to 200 ADP-ribose residues (about 40 residues in plants). The dependence of poly(ADP-ribose) synthesis on DNA strand breaks, and the presence of PARP in multiprotein complexes further containing key effectors of DNA repair, replication and transcription reactions, strongly suggests that this posttranslational modification is involved in metabolism of nucleic acids, and DNA repair. There are also indications that poly (ADP -ribose) synthesis is involved in regulation of cell cycle and cell death.

[4] Poly (ADP-ribosylation) of proteins is transient in living cells. The poly (ADP-ribose) polymers are rapidly turned over, being converted to free ADP-ribose by the exoglycosidase and endoglycosidase activity of poly (ADP-ribose) glycohydrolase (PARG; E.C.3.2.1.143). The most proximal unit of ADP-ribose on the protein acceptor is hydrolysed by the action of another enzyme (ADP-ribosyl protein lyase).

[5] In addition to this positive (DNA-repair associated) effect of PARP on cell survival, there is also a negative effect of PARP. The process of activating PARP upon DNA damage is associated with a rapid lowering of NAD⁺ levels, since each ADP-ribose unit transferred by PARP consumes one molecule of NAD⁺. NAD⁺ depletion in turn results in ATP depletion, because NAD⁺ resynthesis requires at least (depending on the biosynthesis pathway) three molecules of ATP per molecule of NAD⁺. Furthermore, NAD⁺ depletion block glyceraldehyde -3-phosphate dehydrogenase activity, which is required to resynthesize ATP during glycolysis. Finally, NAD⁺ is a key carrier of electrons needed to generate ATP via electron transport and oxidative phosphorylation.

[6] The physiological consequence of NAD⁺ and ATP depletion has been established in the context of DNA-damage induced cell death. It has been shown that the completion of apoptosis is absolutely dependent on the presence of ATP and that, in the absence of this nucleotide, the type of cellular demise switches from apoptosis to necrosis. Since the cellular lysis associates with necrosis generates further damage to neighboring cells it is preferable for multicellular organisms to favour apoptotic cell death rather than necrosis.

[7] It is thus very important to consider the delicate balance of positive and negative effects of the poly (ADP ribosyl)ation on the potential of a cell to survive DNA damage.

[8] WO 00/04173 describes methods to modulate programmed cell death (PCD) in eukaryotic cells and organisms, particularly plant cells and plants, by

introducing of "PCD modulating chimeric genes" influencing the expression and/or apparent activity of endogenous poly-(ADP-ribose) polymerase (PARP) genes. Programmed cell death may be inhibited or provoked. The invention particularly relates to the use of nucleotide sequences encoding proteins with PARP activity for modulating PCD, for enhancing growth rate or for producing stress tolerant cells and organisms.

[9] PARP encoding genes have been identified in a number of animals such as *Rattus norvegicus* (Accession numbers: NM_031339, NW_043030, AB019366,), *Mus musculus* (Accession numbers: NT_039598, NM_003631, AF079557), *Homo sapiens* (Accession numbers: NT_017696; NM_003631, AF005043), *Bos taurus* (Accession numbers: NM_174138, U78975) *Drosophila melanogaster* (Accession number: AF079556)

[10] In plants, a poly(ADP-ribose) glycohydrolase has been identified by map-based cloning of the wild-type gene inactivated in a mutant affected in clock-controlled transcription of genes in *Arabidopsis* and in photoperiod dependent transition from vegetative growth to flowering (*tej*). The nucleotide sequence of the gene can be obtained from nucleotide databases under the accession number AF394690 (Panda et al., 2002 Dev. Cell. 3, 51-61).

Summary of the invention

[11] The invention provides a method to produce a plant tolerant to stress conditions comprising the steps of providing plant cells with a chimeric gene to create transgenic plant cells, wherein the chimeric gene comprises the following operably linked DNA fragments: a plant-expressible promoter; a DNA region, which when transcribed yields an ParG inhibitory RNA molecule; and a 3' end region involved in transcription termination and polyadenylation. A population of transgenic plant lines is regenerated from the transgenic plant cell; and a stress tolerant plant line is identified within the population of transgenic plant lines. The ParG inhibitory RNA molecule may comprise a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present in the plant cell (the endogenous ParG

gene). The ParG inhibitory RNA molecule may also comprise a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the ParG gene present in the plant cell (the endogenous ParG gene). In yet another embodiment, the parG inhibitory RNA may comprise a sense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present in the plant cell and an antisense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the ParG gene present in the plant cell, wherein the sense and antisense region are capable of forming a double stranded RNA region comprising said at least 20 consecutive nucleotides. The chimeric gene may further comprise a DNA region encoding a self-splicing ribozyme between said DNA region coding for parG inhibitory RNA molecule and the 3' end region. Stress conditions may be selected from heat, drought, nutrient depletion, oxidative stress or high light conditions.

[12] In another embodiment of the invention, a method is provided to produce a plant tolerant to stress conditions comprising the steps of: isolating a DNA fragment of at least 100 bp comprising a part of the parG encoding gene of the plant of interest; producing a chimeric gene by operably linking a plant expressible promoter region to the isolated DNA fragment comprising part of the parG encoding gene of the plant in direct orientation compared to the promoter region; and to the isolated DNA fragment comprising part of the parG encoding gene of said plant in inverted orientation compared to the promoter region, and a 3' end region involved in transcription termination and polyadenylation. These chimeric genes are then provided to plant cells to create transgenic plant cells. A population of transgenic plant lines is regenerated from the transgenic plant cells; and a stress tolerant plant line is identified within the population of transgenic plant lines. The invention also relates to stress tolerant plant cells and plants obtained by this process.

[13] The invention also provides DNA molecules comprising a plant-expressible promoter, operably linked to a DNA region, which when transcribed yields an ParG inhibitory RNA molecule, and to a 3' end region

involved in transcription termination and polyadenylation. The ParG inhibitory RNA molecule may comprise a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present in the plant cell (the endogenous ParG gene). The ParG inhibitory RNA molecule may also comprise a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the ParG gene present in the plant cell (the endogenous ParG gene). In yet another embodiment, the parG inhibitory RNA may comprise a sense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present in the plant cell and an antisense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the ParG gene present in the plant cell, wherein the sense and antisense region are capable of forming a double stranded RNA region comprising said at least 20 consecutive nucleotides. The chimeric gene may further comprise a DNA region encoding a self-splicing ribozyme between said DNA region coding for parG inhibitory RNA molecule and the 3' end region.

[14] In yet another embodiment, the invention relates to plant cell comprising the DNA molecule of the invention and plants consisting essentially of such plant cells, as well as to processes for producing stress tolerant plants, comprising the step of further crossing such plants with another plant. Seeds and propagating material of such plants comprising the chimeric genes of the invention are also provided.

Brief description of the figures.

[15] Figure 1. Schematic representation of the poly-ADP ribose polymerization / depolymerization cycle by the action of PARP/PARG in a eukaryotic cell.

[16] Figure 2. Diagram of the NAD⁺ and ATP content of *Arabidopsis* lines under high light stress. Dark boxes represent NAD content under high light conditions expressed as percentage of the value for NAD content determined

under low light conditions. Light boxes represent ATP content under high light conditions expressed as percentage of the value for ATP content determined under low light conditions.

[17] Figure 3. Diagram of the NAD⁺ and ATP content of corn lines under nitrogen depletion stress. Dark boxes represent NAD content while light boxes represent ATP content.

Detailed description of preferred embodiments

[18] The invention is based, on the one hand, on the demonstration that cells from stress resistant plant lines comprising a chimeric gene reducing the PARP gene expression, exhibited a higher NAD/ATP content under adverse conditions than cells from untransformed plant lines. On the other hand, it has been observed that silencing of the expression of PARG encoding gene in tobacco using a transient silencing RNA vector based on satellite viruses resulted in a similar phenotype as that observed for silencing of PARP encoding gene using the same silencing system.

[19] Although not intending to limit the invention to a specific mode of action, it is expected that silencing of PARG gene expression results in a similar phenotype as silencing of PARP gene expression for the following reasons. As can be seen from Figure 1, polymerization of ADP ribose catalyzed by PARP, consuming NAD, is followed by depolymerization of poly ADP ribose, catalyzed by PARG. Poly ADP ribosylation of the PARP protein itself results in inactivation of the PARP protein. The speed at which the ADP ribose polymerization / depolymerization cycle occurs in plant cells, leading to NAD depletion and consequently ATP depletion, can be slowed down or stopped by reduction of the PARP gene expression or of the enzymatic activity of PARP. As a result, plant cells, and plants comprising such cells are more resistant to adverse conditions. The data provided here indicate that a similar effect can be obtained through slowing down or stopping the cycle by reduction of the PARG gene expression or PARG activity.

[20] The invention relates to reduction of plant cell death in response to adverse environmental conditions, and consequently to enhanced stress resistance, by altering the level of expression of ParG genes, or by altering the activity or the apparent activity of PARG proteins in that plant cell. Conveniently, the level of expression of ParG genes may be controlled genetically by introduction of chimeric genes altering the expression of ParG genes, or by altering the endogenous PARG encoding genes.

[21] In one embodiment of the invention, a method for producing plants tolerant to stress conditions or adverse growing conditions is provided comprising the steps of:

- providing plant cells with a chimeric gene to create transgenic plant cells, wherein the chimeric gene comprises the following operably linked DNA fragments:
 - a plant-expressible promoter;
 - a DNA region, which when transcribed yields a ParG inhibitory RNA molecule;
 - a 3' end region involved in transcription termination and polyadenylation;
- regenerating a population of transgenic plant lines from said transgenic plant cell; and
- identifying a stress tolerant plant line within said population of transgenic plant lines.

[22] As used herein "a stress tolerant plant" or "a plant tolerant to stress conditions or adverse growing conditions" is a plant (particularly a plant obtained according to the methods of the invention), which, when subjected to adverse growing conditions for a period of time, such as but not limited to drought, high temperatures, limited supply of nutrients (particularly nitrogen), high light intensities, grows better than a control plant not treated according to the methods of the invention. This will usually be apparent from the general appearance of the plants and may be measured e.g., by increased biomass production, continued vegetative growth under adverse conditions or higher seed yield. Stress tolerant plant have a broader growth spectrum, i.e. they

are able to withstand a broader range of climatological and other abiotic changes, without yield penalty. Biochemically, stress tolerance may be apparent as the higher NAD^+ -NADH /ATP content and lower production of reactive oxygen species of stress tolerant plants compared to control plants under stress condition. Stress tolerance may also be apparent as the higher chlorophyll content, higher photosynthesis and lower chlorophyll fluorescence under stress conditions in stress tolerant plants compared to control plants under the same conditions.

[23] It will be clear that it is also not required that the plant be grown continuously under the adverse conditions for the stress tolerance to become apparent. Usually, the difference in stress tolerance between a plant or plant cell according to the invention and a control plant or plant cell will become apparent even when only a relatively short period of adverse conditions is encountered during growth.

[24] As used herein, a "ParG inhibitory RNA molecule" is an RNA molecule that is capable of decreasing the expression of the endogenous PARG encoding genes of a plant cell, preferably through post-transcriptional silencing. It will be clear that even when a ParG inhibitory RNA molecule decreases the expression of a PARG encoding gene through post-transcriptional silencing, such an RNA molecule may also exert other functions within a cell, such as e.g. guiding DNA methylation of the endogenous ParG gene, again ultimately leading to decreased expression of the PARG encoding gene. Also, expression of the endogenous PARG encoding genes of a plant cell may be reduced by transcriptional silencing, e.g., by using RNAi or dsRNA targeted against the promoter region of the endogenous ParG gene.

[25] As used herein, a "PARG encoding gene" or a "ParG gene" is a gene capable of encoding a PARG (poly ADP ribose glycohydrolase) protein, wherein the PARG protein catalyzes the depolymerization of poly ADP-ribose,

by releasing free ADP ribose units either by endoglycolytic or exoglycolytic action.

[26] PARG encoding genes may comprise a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID No 1 (*Arabidopsis thaliana*) or of SEQ ID No 2 (*Solanum tuberosum*) or parts thereof, such as a DNA fragment comprising the nucleotide sequence of SEQ ID No. 3 or SEQ ID 4.

[27] However, it will be clear that the skilled person can isolate variant DNA sequences from other plant species, by hybridization with a probe derived from the above mentioned PARG encoding genes from plant species, or even with a probe derived from the above mentioned PARG encoding genes from animal species. To this end, the probes should preferably have a nucleotide sequence comprising at least 40 consecutive nucleotides from the coding region of those mentioned PARG encoding genes sequences, preferably from the coding region of SEQ ID No 3 or SEQ ID No 4. The probes may however comprise longer regions of nucleotide sequences derived from the PARG genes, such as about 50, 60, 75, 100, 200 or 500 consecutive nucleotides from any of the mentioned PARG genes. Preferably, the probe should comprise a nucleotide sequence coding for one of the highly conserved regions of the catalytic domain, which have been identified by aligning the different PARG proteins from animals. These regions are also present in the identified PARG protein from *Arabidopsis thaliana* and comprise the amino acid sequence LXVDFANXXXGGG (corresponding to SEQ ID No 1 from the amino acid at position 252 to the amino acid at position 264; X may be any amino acid) LXVDFANXXXGGGXXXXGXVQEEIRF (corresponding to SEQ ID No 1 from the amino acid at position 252 to the amino acid at position 277) or LXVDFANXXXGGGXXXXGXVQEEIRFXXXPE (corresponding to SEQ ID No 1 from the amino acid at position 252 to the amino acid at position 282), TGXWGCXFXGD (corresponding to SEQ ID No 1 from the amino acid at position 449 to the amino acid at position 460) or TGXWGCGAFXGDXXLKXXXQ (corresponding to SEQ ID No 1 from the amino acid at position 449 to the amino acid at position 468). Other conserved

regions have the amino acid sequence DXXRXXXXAIDA (corresponding to SEQ ID No 1 from the amino acid at position 335 to the amino acid at position 344) or REXKAXXGF (corresponding to SEQ ID No 1 from the amino acid at position 360 to the amino acid at position 369) or GXXXXSXYTGY (corresponding to SEQ ID No 1 from the amino acid at position 303 to the amino acid at position 313). Hybridization should preferably be under stringent conditions.

[28] "Stringent hybridization conditions" as used herein mean that hybridization will generally occur if there is at least 95% and preferably at least 97% sequence identity between the probe and the target sequence. Examples of stringent hybridization conditions are overnight incubation in a solution comprising 50% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared carrier DNA such as salmon sperm DNA, followed by washing the hybridization support in 0.1 x SSC at approximately 65 °C. Other hybridization and wash conditions are well known and are exemplified in Sambrook et al, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, NY (1989), particularly chapter 11.

[29] Alternatively, ParG encoding genes or parts thereof may also be isolated by PCR based techniques, using as primers oligonucleotides comprising at least 20 consecutive nucleotides from a nucleotide sequence of the mentioned PARG encoding genes or the complement thereof. Such primers may comprise a nucleotide sequence encoding a conserved region, as mentioned above, or be complementary to such a nucleotide sequence. Oligonucleotides which may be used for that purpose may comprise the nucleotide sequence of either or SEQ ID No.5, SEQ ID No 6., SEQ ID No. 7 or SEQ ID No. 8.

[30] Specific PCR fragments from ParG genes may e.g., be obtained by using combinations of the oligonucleotides having the nucleotide sequence of SEQ ID No. 5 and SEQ ID No 6 using e.g., *Arabidopsis* genomic DNA or

cDNA as a template DNA, or by using combinations of the oligonucleotides having the nucleotide sequence of SEQ ID No. 7 and SEQ ID No 8 using e.g., potato genomic DNA or cDNA as a template DNA, under stringent annealing conditions.

[31] The isolated sequences may encode a functional PARG protein or a part thereof. Preferably the isolated sequences should comprise a nucleotide sequence coding for one or more of the highly conserved regions from the catalytic domain of PARG proteins as mentioned elsewhere.

[32] However, for the purpose of the invention is not required that the isolated sequences encode a functional ParG protein nor that a complete coding region is isolated. Indeed, all that is required for the invention is that a chimeric gene can be designed or produced, based on the identified or isolated sequence of the endogenous ParG gene from a plant, which is capable of producing a ParG inhibitory RNA. Several alternative methods are available to produce such a ParG inhibitory RNA molecule.

[33] In one embodiment, the ParG inhibitory RNA molecule encoding chimeric gene is based on the so-called antisense technology. In other words, the coding region of the chimeric gene comprises a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the endogenous ParG gene of the plant cell or plant, the expression of which is targeted to be reduced. Such a chimeric gene may be conveniently constructed by operably linking a DNA fragment comprising at least 20 nucleotides from the isolated or identified ParG gene, or part of such a gene, in inverse orientation, to a plant expressible promoter and 3'end formation region involved in transcription termination and polyadenylation. It will be immediately clear that there is no need to know the exact nucleotide sequence or complete nucleotide sequence of such a DNA fragment from an isolated ParG gene.

[34] In another embodiment the ParG inhibitory RNA molecule encoding chimeric gene is based on the so-called co-suppression technology. In other

words, the coding region of the chimeric gene comprises a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the endogenous ParG gene of the plant cell or plant, the expression of which is targeted to be reduced. Such a chimeric gene may be conveniently constructed by operably linking a DNA fragment comprising at least 20 nucleotides from the isolated or identified ParG gene, or part of such a gene, in direct orientation, to a plant expressible promoter and 3'end formation region involved in transcription termination and polyadenylation. Again it is not required to know the exact nucleotide sequence of the used DNA fragment from the isolated ParG gene.

[35] The efficiency of the above mentioned chimeric genes in reducing the expression of the endogenous ParG gene may be further enhanced by inclusion of DNA elements which result in the expression of aberrant, unpolyadenylated ParG inhibitory RNA molecules. One such DNA element suitable for that purpose is a DNA region encoding a self-splicing ribozyme, as described in WO 00/01133.

[36] The efficiency of the above mentioned chimeric genes in reducing the expression of the endogenous ParG gene of a plant cell may also be further enhanced by including into one plant cell simultaneously a chimeric gene as herein described encoding an antisense ParG inhibitory RNA molecule and a chimeric gene as herein described encoding a sense ParG inhibitory RNA molecule, wherein said antisense and sense ParG inhibitory RNA molecules are capable of forming a double stranded RNA region by base pairing between the mentioned at least 20 consecutive nucleotides, as described in WO 99/53050.

[37] As further described in WO 99/53050, the sense and antisense ParG inhibitory RNA regions, capable of forming a double stranded RNA region may be present in one RNA molecule, preferably separated by a spacer region. The spacer region may comprise an intron sequence. Such a chimeric gene may be conveniently constructed by operably linking a DNA fragment comprising at least 20 nucleotides from the isolated or identified endogenous

ParG gene, the expression of which is targeted to be reduced, in an inverted repeat, to a plant expressible promoter and 3' end formation region involved in transcription termination and polyadenylation. To achieve the construction of such a chimeric gene, use can be made of the vector described in WO 02/059294

[38] An embodiment of the invention thus concerns a method for obtaining a stress tolerant plant line comprising the steps of

- providing plant cells with a chimeric gene to create transgenic plant cells, wherein the chimeric gene comprises the following operably linked DNA fragments:

- a plant-expressible promoter;
- a DNA region, which when transcribed yields a ParG inhibitory RNA molecule comprising a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present in said plant cell; or
- a DNA region, which when transcribed yields a ParG inhibitory RNA molecule comprising a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the ParG gene present in said plant cell; or
- a DNA region, which when transcribed yields a ParG inhibitory RNA molecule comprising a sense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present in said plant cell and an antisense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the ParG gene present in said plant cell, wherein said sense and antisense region are capable of forming a double stranded RNA region comprising said at least 20 consecutive nucleotides.
- a 3' end region involved in transcription termination and polyadenylation;
- regenerating a population of transgenic plant lines from said transgenic plant cell; and

- identifying a stress tolerant plant line within said population of transgenic plant lines.

[39] As used herein "comprising" is to be interpreted as specifying the presence of the stated features, integers, steps or components as referred to, but does not preclude the presence or addition of one or more features, integers, steps or components, or groups thereof. Thus, e.g., a nucleic acid or protein comprising a sequence of nucleotides or amino acids, may comprise more nucleotides or amino acids than the actually cited ones, i.e., be embedded in a larger nucleic acid or protein. A chimeric gene comprising a DNA region which is functionally or structurally defined, may comprise additional DNA regions etc.

[40] It will thus be clear that the minimum nucleotide sequence of the antisense or sense RNA region of about 20 nt of the ParG coding region may be comprised within a larger RNA molecule, varying in size from 20 nt to a length equal to the size of the target gene.

[41] The mentioned antisense or sense nucleotide regions may thus be about from about 21 nt to about 5000 nt long, such as 21nt, 40 nt, 50 nt, 100nt, 200 nt, 300nt, 500nt, 1000 nt, 2000 nt or even about 5000 nt or larger in length.

[42] Moreover, it is not required for the purpose of the invention that the nucleotide sequence of the used inhibitory ParG RNA molecule or the encoding region of the chimeric gene, is completely identical or complementary to the endogenous ParG gene the expression of which is targeted to be reduced in the plant cell. The longer the sequence, the less stringent the requirement for the overall sequence identity is. Thus, the sense or antisense regions may have an overall sequence identity of about 40 % or 50% or 60 % or 70% or 80% or 90 % or 100% to the nucleotide sequence of the endogenous ParG gene or the complement thereof. However, as mentioned antisense or sense regions should comprise a nucleotide sequence of 20 consecutive nucleotides having about 100% sequence identity

to the nucleotide sequence of the endogenous ParG gene. Preferably the stretch of about 100 % sequence identity should be about 50, 75 or 100 nt.

[43] For the purpose of this invention, the "sequence identity" of two related nucleotide sequences, expressed as a percentage, refers to the number of positions in the two optimally aligned sequences which have identical residues (x100) divided by the number of positions compared. A gap, i.e. a position in an alignment where a residue is present in one sequence but not in the other is regarded as a position with non-identical residues. The alignment of the two sequences is performed by the Needleman and Wunsch algorithm (Needleman and Wunsch 1970) Computer-assisted sequence alignment, can be conveniently performed using standard software program such as GAP which is part of the Wisconsin Package Version 10.1 (Genetics Computer Group, Madison, Wisconsin, USA) using the default scoring matrix with a gap creation penalty of 50 and a gap extension penalty of 3.

[44] It will be clear that whenever nucleotide sequences of RNA molecules are defined by reference to nucleotide sequence of corresponding DNA molecules, the thymine (T) in the nucleotide sequence should be replaced by uracil (U). Whether reference is made to RNA or DNA molecules will be clear from the context of the application.

[45] It will also be clear that chimeric genes capable of producing inhibitory ParG genes for a particular ParG gene in a particular plant variety or plant species, may also be used to inhibit ParG gene expression in other plant varieties or plant species. Indeed, when sufficient homology exists between the ParG inhibitory RNA region and the ParG gene, or when the ParG genes share the same stretch of 19 nucleotides, expression of those other genes will also be down-regulated.

[46] In view of the potential role of ParG in nucleic acid metabolism, it may be advantageous that the expression of the endogenous ParG gene by the ParG inhibitory RNA is not completely inhibited. Downregulating the expression of a particular gene by gene silencing through the introduction of a

chimeric gene encoding ParG inhibitory RNA will result in a population of different transgenic lines, exhibiting a distribution of different degrees of silencing of the ParG gene. The population will thus contain individual transgenic plant lines, wherein the endogenous ParG gene is silenced to the required degree of silencing. A person skilled in the art can easily identify such plant lines, e.g. by subjecting the plant lines to a particular adverse condition, such a high light intensity, oxidative stress, drought, heat etc. and selecting those plants which perform satisfactory and survive best the treatment.

[47] As used herein, the term "promoter" denotes any DNA which is recognized and bound (directly or indirectly) by a DNA-dependent RNA-polymerase during initiation of transcription. A promoter includes the transcription initiation site, and binding sites for transcription initiation factors and RNA polymerase, and can comprise various other sites (e.g., enhancers), at which gene expression regulatory proteins may bind.

[48] The term "regulatory region", as used herein, means any DNA, that is involved in driving transcription and controlling (i.e., regulating) the timing and level of transcription of a given DNA sequence, such as a DNA coding for a protein or polypeptide. For example, a 5' regulatory region (or "promoter region") is a DNA sequence located upstream (i.e., 5') of a coding sequence and which comprises the promoter and the 5'-untranslated leader sequence. A 3' regulatory region is a DNA sequence located downstream (i.e., 3') of the coding sequence and which comprises suitable transcription termination (and/or regulation) signals, including one or more polyadenylation signals.

[49] In one embodiment of the invention the promoter is a constitutive promoter. In another embodiment of the invention, the promoter activity is enhanced by external or internal stimuli (inducible promoter), such as but not limited to hormones, chemical compounds, mechanical impulses, abiotic or biotic stress conditions. The activity of the promoter may also regulated in a temporal or spatial manner (tissue-specific promoters; developmentally regulated promoters).

[50] For the purpose of the invention, the promoter is a plant-expressible promoter. As used herein, the term "plant-expressible promoter" means a DNA sequence which is capable of controlling (initiating) transcription in a plant cell. This includes any promoter of plant origin, but also any promoter of non-plant origin which is capable of directing transcription in a plant cell, i.e., certain promoters of viral or bacterial origin such as the CaMV35S (Hapster et al., 1988), the subterranean clover virus promoter No 4 or No 7 (WO9606932), or T-DNA gene promoters but also tissue-specific or organ-specific promoters including but not limited to seed-specific promoters (e.g., WO89/03887), organ-primordia specific promoters (An et al., 1996), stem-specific promoters (Keller et al., 1988), leaf specific promoters (Hudspeth et al., 1989), mesophyl-specific promoters (such as the light-inducible Rubisco promoters), root-specific promoters (Keller et al., 1989), tuber-specific promoters (Keil et al., 1989), vascular tissue specific promoters (Peleman et al., 1989), stamen-selective promoters (WO 89/10396, WO 92/13956), dehiscence zone specific promoters (WO 97/13865) and the like.

[51] Methods for the introduction of chimeric genes into plants are well known in the art and include *Agrobacterium*-mediated transformation, particle gun delivery, microinjection, electroporation of intact cells, polyethyleneglycol-mediated protoplast transformation, electroporation of protoplasts, liposome-mediated transformation, silicon-whiskers mediated transformation etc. The transformed cells obtained in this way may then be regenerated into mature fertile plants.

[52] The transgenic plant cells and plant lines according to the invention may further comprise chimeric genes which will reduce the expression of PARP genes as described in WO 00/04173. These further chimeric genes may be introduced e.g. by crossing the transgenic plant lines of the current invention with transgenic plants containing PARP gene expression reducing chimeric genes. Transgenic plant cells or plant lines may also be obtained by introducing or transforming the chimeric genes of the invention into transgenic plant cells comprising the PARP gene expression reducing chimeric genes or

vice versa. Alternatively, the PARP and PARG inhibitory RNA regions may be encoded by one chimeric gene and transcribed as one RNA molecule.

[53] The chimeric genes of the invention (or the inhibitory RNA molecules corresponding thereto) may also be introduced into plant cells in a transient manner, e.g. using the viral vectors, such as viral RNA vectors as described in WO 00/63397 or WO 02/13964.

[54] Having read this specification, it will be immediately clear to the skilled artisan, that mutant plant cells and plant lines, wherein the PARG activity is reduced may be used to the same effect as the transgenic plant cells and plant lines described herein. Mutants in ParG gene of a plant cell or plant may be easily identified using screening methods known in the art, whereby chemical mutagenesis, such as e.g., EMS mutagenesis, is combined with sensitive detection methods (such as e.g., denaturing HPLC). An example of such a technique is the so-called "Targeted Induced Local Lesions in Genomes" method as described in McCallum et al, Plant Physiology 123 439-442 or WO 01/75167. However, other methods to detect mutations in particular genome regions or even alleles, are also available and include screening of libraries of existing or newly generated insertion mutant plant lines, whereby pools of genomic DNA of these mutant plant lines are subjected to PCR amplification using primers specific for the inserted DNA fragment and primers specific for the genomic region or allele, wherein the insertion is expected (see e.g. Maes et al., 1999, Trends in Plant Science, 4, pp 90-96).

[55] Plant cell lines and plant lines may also be subjected to mutagenesis by selection for resistance to ParG inhibitors, such as gallotannins . (Ying, et al. (2001). Proc. Natl. Acad. Sci. USA 98(21), 12227-12232; Ying, W., Swanson, R.A. (2000). NeuroReport 11(7), 1385-1388.

[56] Thus, methods are available in the art to identify plant cells and plant lines comprising a mutation in the ParG gene. This population of mutant cells or plant lines can then be subjected to different abiotic stresses, and their

phenotype or survival can be easily determined. Additionally, the NAD and/or the ATP content of the stressed cells can be determined and compared to results of such determinations of unstressed cells. In stress tolerant cells, the reduction of NAD content under stress conditions should when compared with unstressed cells, should be lower than for corresponding control cells.

[57] It is also an object of the invention to provide plant cells and plants containing the chimeric genes or the RNA molecules according to the invention. Gametes, seeds, embryos, either zygotic or somatic, progeny or hybrids of plants comprising the chimeric genes of the present invention, which are produced by traditional breeding methods are also included within the scope of the present invention.

[58] The plants obtained by the methods described herein may be further crossed by traditional breeding techniques with other plants to obtain stress tolerant progeny plants comprising the chimeric genes of the present invention.

[59] The methods and means described herein are believed to be suitable for all plant cells and plants, both dicotyledonous and monocotyledonous plant cells and plants including but not limited to cotton, Brassica vegetables, oilseed rape, wheat, corn or maize, barley, alfalfa, peanuts, sunflowers, rice, oats, sugarcane, soybean, turf grasses, barley, rye, sorghum, sugar cane, vegetables (including chicory, lettuce, tomato, zucchini, bell pepper, eggplant, cucumber, melon, onion, leek), tobacco, potato, sugarbeet, papaya, pineapple, mango, *Arabidopsis thaliana*, but also plants used in horticulture, floriculture or forestry (poplar, fir, eucalyptus etc.).

[60] The following non-limiting Examples describe method and means for increasing stress tolerance in plants according to the invention.

[61] Unless stated otherwise in the Examples, all recombinant DNA techniques are carried out according to standard protocols as described in Sambrook et al. (1989) Molecular Cloning: A Laboratory Manual, Second

Edition, Cold Spring Harbor Laboratory Press, NY and in Volumes 1 and 2 of Ausubel et al. (1994) Current Protocols in Molecular Biology, Current Protocols, USA. Standard materials and methods for plant molecular work are described in Plant Molecular Biology Labfax (1993) by R.D.D. Croy, jointly published by BIOS Scientific Publications Ltd (UK) and Blackwell Scientific Publications, UK. Other references for standard molecular biology techniques include Sambrook and Russell (2001) Molecular Cloning: A Laboratory Manual, Third Edition, Cold Spring Harbor Laboratory Press, NY, Volumes I and II of Brown (1998) Molecular Biology LabFax, Second Edition, Academic Press (UK). Standard materials and methods for polymerase chain reactions can be found in Dieffenbach and Dveksler (1995) PCR Primer: A Laboratory Manual, Cold Spring Harbor Laboratory Press, and in McPherson et al. (2000) PCR - Basics: From Background to Bench, First Edition, Springer Verlag, Germany.

[62] Throughout the description and Examples, reference is made to the following sequences:

- [63] SEQ ID N°1: amino acid sequence of the ParG protein from *Arabidopsis thaliana*.
- [64] SEQ ID N°2: amino acid sequence of part of the ParG protein from *Solanum tuberosum*.
- [65] SEQ ID N°3: nucleotide sequence encoding the ParG protein from *Arabidopsis thaliana*.
- [66] SEQ ID N°4: nucleotide sequence encoding the part of the ParG protein from *Solanum tuberosum*.
- [67] SEQ ID N°5: nucleotide sequence of an oligonucleotide primer suitable for PCR amplification of part of a ParG protein encoding DNA fragment.
- [68] SEQ ID N°6: nucleotide sequence of an oligonucleotide primer suitable for PCR amplification of part of a ParG protein encoding DNA fragment.
- [69] SEQ ID N°7: nucleotide sequence of an oligonucleotide primer suitable for PCR amplification of part of a ParG protein encoding DNA fragment.
- [70] SEQ ID N°8: nucleotide sequence of an oligonucleotide primer suitable for PCR amplification of part of a ParG protein encoding DNA fragment.

- [71] SEQ ID N°9: nucleotide sequence of the T-DNA vector containing the ParG expression reducing chimeric gene based on the *Arabidopsis* ParG gene sequence.
- [72] SEQ ID N°10: amino acid sequence of conserved sequence 1 of PARG proteins.
- [73] SEQ ID N°11: amino acid sequence of conserved sequence 2 of PARG proteins.
- [74] SEQ ID N°12: amino acid sequence of conserved sequence 3 of PARG proteins.
- [75] SEQ ID N°13: amino acid sequence of conserved sequence 4 of PARG proteins.
- [76] SEQ ID N°14: amino acid sequence of conserved sequence 5 of PARG proteins.

Examples

Example 1. Analysis of the influence of stress on energy production efficiency of transgenic stress tolerant plant lines containing PARP gene expression reducing chimeric genes.

- [77] Hypocotyls of transgenic *Brassica napus* plants comprising PARP gene expression reducing chimeric genes as described in WO 00/04173 were cultivated for 5 days on a growth medium. Explants were then transferred to liquid medium comprising 30 mg/L aspirin or acetylsalicylic acid (resulting in oxidative stress conditions) for one day. In control experiments, hypocotyls of non-transgenic *Brassica napus* plants N90-740 were cultivated on the same growth medium and then incubated for one day in liquid medium comprising 30 mg/L aspirin. In addition, hypocotyls of both the transgenic lines and the control line were cultivated on the same growth medium without aspirin.
- [78] After the cultivation period, the ATP content of 125 explants was determined for each experiment. Additionally, the oxygen consumed in 3 hours by 125 explants was determined. The results are summarized in Table 1. The standard error of the mean was less than 6%. Whereas, the ratio of moles ATP per mg consumed oxygen in the control plants decreased in the control plants when oxidative stress was applied, the same ratio in the stress tolerant transgenic plant lines actually increased under stress conditions, and was considerably higher (about 24%) than in the control plants. The stress resistant transgenic lines thus maintained an constant energy production efficiency, whereas the control lines exhibited an decreased energy production efficiency. In addition, superoxide production, expressed as a percentage of superoxide production in control plants not subjected to the oxidative stress, did not increase in stress tolerant plants subjected to stress conditions.

[79] Table 1. Influence of stress on energy production efficiency of 5 days cultured *Brassica napus* hypocotyl explants.

Plant line	Stress	moles ATP per 125 explants	O ₂ mg/L consumed in 3 hrs by 125 explants	moles ATP	Superoxide production
				mg consumed O ₂	
N90-740 (control)	None	12.4×10^{-7}	2.96	4.19×10^{-7}	100%
	30mg/L aspirin	13.2×10^{-7}	4.06	3.25×10^{-7}	167%
Transgenic line	None	9.3×10^{-7}	2.33	3.99×10^{-7}	108%
	30mg/L aspirin	11.4×10^{-7}	2.82	4.04×10^{-7}	100%

[80] In another experiment, the NAD⁺ and ATP content of 4 different transgenic *Arabidopsis* lines comprising PARP gene expression reducing chimeric genes as described in WO 00/04173 were determined under high and low light conditions, and compared to the values obtained for a non transformed control line under the same conditions. The 4 different lines exhibited different degrees of stress resistance as exhibited e.g. by their ability to withstand heat and/or drought conditions. The values obtained for the NAD and ATP contents under high light stress are expressed as a percentage of the values for the NAD and ATP contents under low light conditions, and are plotted in Figure 2.

[81] The results show that high light stress leads to a significant NAD reduction in control plant cells and in the transgenic plant line which is the least stress resistant. The more stress resistant the transgenic plant lines are, the less significant the NAD reduction is under high light stress conditions.

[82] In another experiment, the NAD⁺ and ATP content of a segregating population resulting from a cross between transgenic corn lines comprising PARP gene expression reducing chimeric genes as described in WO 00/04173 and an untransformed corn line, were determined under conditions of nutrient (nitrogen) depletion, and compared to the values obtained for a non transformed control line under the same conditions. Figure 3 is a graphic representation of the of the obtained results. Hemizygous and azygous lines were discriminated by verification for the presence of the selectable marker gene. The NAD and ATP content was significantly higher in the hemizygous, stress tolerant plants than in the untransformed control plants or the azygous plants.

Example 2. Construction of ParG gene expression reducing chimeric genes.

- [83] To reduce the expression of the PARG gene e.g. in *Arabidopsis* and related plants, a chimeric gene was constructed which is capable expressing a dsRNA comprising both a sense and antisense region which can form a double stranded RNA. Such dsRNA is very effective in reducing the expression of the genes with which is shares sequence homology, by post-transcriptional silencing. The chimeric gene comprises the following DNA fragments:
- A promoter region from Cauliflower mosaic Virus (CaMV 35S);
 - A DNA fragment comprising 163 bp from the ParG gene from *Arabidopsis thaliana* in direct orientation (Genbank Accession number AF394690 from nucleotide position 973 to 1135);
 - A DNA fragment encoding intron 2 from the pdk gene from *Flaveria*;
 - The DNA fragment comprising 163 bp from the ParG gene from *Arabidopsis thaliana* in inverted orientation (Genbank Accession number AF394690 from nucleotide position 973 to 1135)
 - A fragment of the 3' untranslated end from the octopine synthetase gene from *Agrobacterium tumefaciens*.

- [84] This chimeric gene was introduced in a T-DNA vector, between the left and right border sequences from the T-DNA, together with a selectable marker gene providing resistance to the herbicide phosphinotricin.
- [85] To reduce the expression of the PARC gene e.g. in potatoes and related plants, a chimeric gene is constructed which is capable expressing a dsRNA comprising both a sense and antisense region of a cDNA sequence from potato, that is capable of encoding a protein having high sequence identity with the N-terminal part of the *Arabidopsis* PARC protein. The chimeric gene comprises the following DNA fragments:
- A promoter region from Cauliflower mosaic Virus (CaMV 35S);
 - A DNA fragment comprising a sequence of at least 100 bp from ParG homologue from *Solanum tuberosum* in direct orientation (Genbank Accession number BE340510);
 - A DNA fragment encoding intron 2 from the pdk gene from *Flaveria*;
 - The DNA fragment comprising the sequence of at least 100 bp from ParG homologue from *Solanum tuberosum* in inverted orientation (Genbank Accession number BE340510);
 - A fragment of the 3' untranslated end from the octopine synthetase gene from *Agrobacterium tumefaciens*

- [86] This chimeric gene is introduced in a T-DNA vector, between the left and right border sequences from the T-DNA, together with a selectable marker gene providing resistance to the herbicide phosphinotricin.

Example 3. Analysis of transgenic plant lines comprising ParG gene expression reducing chimeric genes.

- [87] The chimeric genes of Example 2 are introduced into *Arabidopsis* or potato respectively, by *Agrobacterium* mediated transformation.

[88] The population of obtained transgenic lines is subjected to the following stress conditions, together with control plants:

- Increased heat for a period of days (greenhouse) or hours (in vitro)
- Drought for a period of days
- High light conditions for a period of days
- Nutrient depletion

[89] Individual plant lines surviving well the above mentioned stress conditions are selected.

[90] The NAD content and ATP content for the above mentioned plants is determined under control and stress conditions.

Example 4. Quantitative determination of NAD, ATP and superoxide radicals in plant cells.

[91] Quantification of ATP in plant tissues was done basically as described by Rawlyer et al. (1999), Plant Physiol. 120, 293-300. The assay was used for the determination of the ATP content of hypocotyl explants that were cultured for 4-5 days on A2S3 medium or 2 weeks old *in vitro* cultured *Arabidopsis* plants. All manipulations are performed on crushed ice unless otherwise indicated.

[92] ATP extraction

- Freeze plant material with liquid nitrogen
 - 100 hypocotyl explants
 - \pm 700mg *Arabidopsis* plants (roots + shoots) (about 32-37 18-days old C24 plants)
- Put frozen hypocotyls in mortar and add 6ml of 6% perchloric acid.
- Extraction can be done at room temperature using a pestle. After extraction, put samples as soon as possible on ice.
- Centrifuge at 24,000g (Sorvall, SS34 rotor at 14,000rpm) for 10min. at

4°C.

- The supernatant is neutralized with 5M K₂CO₃ (add 350µl of 5M K₂CO₃ to 3ml of supernatant).
- KClO₄ is removed by spinning as described above.

[93] Quantitative bioluminescent determination of ATP

- The ATP bioluminescent assay kit from Sigma is used (FL-AA).
- Dilute extract 6000 x (about 6 mL extract from which 100µl is taken, that is diluted 1000 times) The dilutions are made with the 'ATP assay mix dilution buffer' (FL-AAB) of the ATP bioluminescent assay kit
- The amount of light that is produced is measured with the TD-20/20 luminometer of Turner Designs (Sunnyvale, USA).
- Standard curve: dissolve ATP standard of kit (FL-AAS) in 10ml of water (2x10⁻⁶ moles)

[94] Quantification of NAD⁺ and NADH in plant tissues was performed, essentially as described by Karp et al. (1983) or Fillpovic et al. (1999) on the following plant material:

Brassica napus: 150 5-days cultured hypocotyl explants/sample

Arabidopsis: 1000mg 18-days old in vitro grown plants (shoots + roots)/sample (corresponds to ±60 C24 plants)

[95] Assay solution.

- (A) For measuring NADH: 25mM potassium phosphate buffer pH7
0.1mM DTT
3µM FMN (Fluka, 83810)
30µM n-decanal (Sigma, D-7384)

(B) For measuring NAD⁺ + NADH:

idem as for measuring NADH alone
+ 2µg/mL alcohol dehydrogenase (Roche, 102 717)

[96] Extraction

- Freeze with liquid nitrogen
- Put frozen plant material in cooled mortar (cooled at -20°C) and add 5mL extraction buffer
- Grind material using a pestle
- Centrifuge at 24 000g (Sorvall, SS34 rotor at 14 000rpm) for 15 minutes at 4°C
- Take 1 mL of supernatant for analysis

[97] Assay***NADH***

- 390 μL of assay solution A
- + 10 μL extract
- + 2 μL NAD(P)H:FMN oxidoreductase
- + 100 μL luciferase solution

NAD⁺ + NADH

- 390 μL of assay solution B
- + 10 μL extract
- 2 minutes at room temperature
- + 2 μL NAD(P)H:FMN oxidoreductase
- + 100 μL luciferase solution

The amount of light that is produced is measured with the TD-20/20 luminometer of Turner Designs (Sunnyvale, USA)

NADH-standard

NADH stock solution: 1mM (7.1mg/10mL H_2O)

NADH: disodium salt, Roche, 107 735

Dilution series in 10mM potassium phosphate buffer pH7: (10^{-2}); 5×10^{-3} ; 2×10^{-3} ; 10^{-3} ; 5×10^{-4}

Add 10 μL of dilutions in 390 μL of assay solution A and perform reaction

Make standard curve

- [98]** Superoxide radicals production was measured by quantifying the reduction of XTT as described in De Block and De Brouwer (2002) Plant Physiol. Biochem. 40, 845-852

[99] BRASSICA NAPUS**[100] Media and reaction buffers****Sowing medium (medium 201):**

Half concentrated Murashige and Skoog salts

2% sucrose

pH 5.8

0.6% agar (Difco Bacto Agar)

250mg/l triacillin

Callus inducing medium A2S3:

MS medium, 0.5g/l Mes (pH 5.8), 3% sucrose, 40mg/l adenine-SO₄,
0.5% agarose, 1mg/l 2,4-D, 0.25mg/l NAA, 1mg/l BAP, 250mg/l
triacillin

Incubation medium:

25mM K-phosphate buffer pH5.8

2% sucrose

1 drop Tween20 for 25ml medium

Reaction buffer:

50mM K-phosphate buffer pH7.4

1mM sodium,3'-{1-[phenylamino-carbonyl]-3,4-tetrazolium}-bis(4-methoxy-6-nitro) = XTT (bts, Germany, cat n° 2525)

1 drop Tween20 for 25ml buffer

- [101] Sterilization of seeds - pregermination of seeds - growing of the seedlings.** Seeds are soaked in 70% ethanol for 2 min, then surface-sterilized for 15 min in a sodium hypochlorite solution (with about 6% active chlorine) containing 0.1% Tween20. Finally, the seeds are rinsed with 1l of sterile tap water. Incubate seeds for at least one hour in sterile tap water (to allow diffusion from seeds of components that may inhibit germination). Seeds are put in 250ml erlenmeyer flasks containing 50ml of sterile tap water (+ 250mg/l triacillin). Shake for

about 20 hours. Seeds from which the radicle is protruded are put in Vitro Vent containers from Duchefa containing about 125ml of sowing medium (10 seeds/vessel, not too many to reduce loss of seed by contamination). The seeds are germinated at $\pm 24^{\circ}\text{C}$ and 10-30:Einsteins $\text{s}^{-1}\text{m}^{-2}$ with a daylength of 16h.

[102] Preculture of the hypocotyl explants and induction of stress

- 12-14 days after sowing, the hypocotyls are cut in about 7-10mm segments.
- The hypocotyl explants (25 hypocotyls/Optilux Petridish, Falcon S1005, Denmark) are cultured for 5 days on medium A2S3 at 25°C (at 10-30 Einsteins $\text{s}^{-1}\text{m}^{-2}$).

[103] XTT-assay

- Transfer 150 hypocotyl explants to a 50ml Falcon tube.
- Wash with reaction buffer (without XTT).
- Add 20mL reaction buffer + XTT.
(explants have to be submerged, but do not vacuum infiltrate)
- Incubate in the dark at 26°C for about 3 hours
- Measure the absorption of the reaction medium at 470nm

[104] ARABIDOPSIS THALIANA

[105] Media and reaction buffers

Plant medium:

Half concentrated Murashige and Skoog salts
B5 vitamins
1.5% sucrose
pH 5.8
0.7% Difco agar

Incubation medium:

10mM K-phosphate buffer pH5.8
2% sucrose
1 drop Tween20 for 25ml medium

Reaction buffer:

50mM K-phosphate buffer pH7.4

1mM sodium,3'-{1-[phenylamino-carbonyl]-3,4-tetrazolium}-bis(4-methoxy-6-nitro) = XTT (bts, Germany, cat n° 2525)

1 drop Tween20 for 25ml buffer

[106] *Arabidopsis* plants

- *Arabidopsis* lines: control

lines to test

- Sterilization of *Arabidopsis* seeds:

2min. 70% ethanol

10 min. bleach (6% active chlorine) + 1drop Tween 20 for 20ml solution

wash 5 times with sterile tap water

- Pregermination of seeds:

In 9cm Optilux Petridishes (Falcon) containing 12ml sterile tap water.

Low light overnight to 24 hours.

- Growing of *Arabidopsis* plants

Seeds are sown in Intergrid Tissue Culture disks of Falcon (nr. 3025) containing ± 125 ml of plant medium: 1 seed/grid.

Plants are grown at 24°C

$30 \mu\text{Einstein s}^{-1}\text{m}^{-2}$

16hours light - 8hours dark

for about 3 weeks (before bolting)

[107] XTT-assay

Control condition (no stress)

- Harvest shoots (roots included) from agar plates and put them directly in a 50ml Falcon tube containing reaction buffer (without XTT)

Stressed shoots

- Transfer shoots to 50ml Falcon tubes containing reaction buffer (without XTT)
- Replace reaction buffer with buffer containing XTT (40mL/tube)

- Shoots have to be submerged, but do not vacuum infiltrate
- Incubate in the dark at 26°C for about 3 hours
- Measure the absorption of the reaction medium at 470nm

[108] Quantification of respiration by measuring oxygen consumption using a Clark polarographic electrode was done in the following way:

[109] Plant material

Brassica napus

150-200* hypocotyl explants

Cultured for 5 days at 25°C

(cfr. protocol vigour assay)

* 150 explants error <10%; 200 explants error <6%

Arabidopsis

For C24 ± 1000mg* *in vitro* plants (shoots + roots) (corresponds with ~50 18-days old plants)

Pregerminate seeds before sowing

Grow for 18 days at 24°C

(cfr. protocol *in vitro* growth Arabidopsis)

* for error <8%

[110] Incubation media

Brassica napus

25mM K-phosphate buffer pH5.8

2% sucrose

Tween20 (1 drop/25ml)

Arabidopsis

10mM K-phosphate buffer pH5.8

2% sucrose

[111] Assay

- Put explants in 100ml glass bottle (Schott, Germany) filled with incubation medium. Put the same weight of shoots in each bottle (\pm 700mg)
- Fill bottle to overflowing and close tightly (avoid large air bubbles)
- Fill also a bottle with incubation medium that does not contain explants (blanco)
- Incubate at 24°C at low light for: 3–4 hours (*Brassica napus*)
 3 hours (*Arabidopsis*)
- Shake gently during incubation (to avoid oxygen depletion of medium around explants)
- Measure oxygen concentration (mg/l) of incubation media using an hand-held dissolved oxygen meter (Cyberscan DO 310; Eutech Instruments, Singapore)
- mg/l consumed oxygen = [oxygen] blanco – [oxygen] sample.

1. A method to produce a plant tolerant to stress conditions comprising the steps of

(a) providing plant cells with a chimeric gene to create transgenic plant cells, said chimeric gene comprising the following operably linked DNA fragments

- (i) a plant-expressible promoter;
- (ii) a DNA region, which when transcribed yields an ParG inhibitory RNA molecule;
- (iii) a 3' end region involved in transcription termination and polyadenylation;

(b) regenerating a population of transgenic plant lines from said transgenic plant cell; and

(c) identifying a stress tolerant plant line within said population of transgenic plant lines.

2. The method according to claim 1, wherein said parG inhibitory RNA molecule comprises comprising a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present in said plant cell.

3. The method according to claim 1, wherein said parG inhibitory RNA molecule is comprises comprising a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the ParG gene present in said plant cell.

4. The method according to claim 2 or 3, wherein said chimeric gene further comprises a DNA region encoding a self-splicing ribozyme between said DNA region coding for said parG inhibitory RNA molecule and said 3' end region.

5. The method according to claim 1, wherein said parG inhibitory RNA comprises a sense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present in said plant cell and an antisense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide

sequence of the ParG gene present in said plant cell, wherein said sense and antisense region are capable of forming a double stranded RNA region comprising said at least 20 consecutive nucleotides.

6. The method according to any one of claims 1 to 5 wherein said stress conditions is selected from heat, drought, nutrient depletion, oxidative stress or high light conditions.

7. The method according to any one of claims 1 to 6, comprising further crossing said transgenic plant line with another plant line to obtain stress tolerant progeny plants.

8. A method to produce a plant tolerant to stress conditions comprising the steps of:

(a) isolating a DNA fragment of at least 100 bp comprising a part of the parG encoding gene of said plant;

(b) producing a chimeric gene by operably linking the following DNA fragments;

(i) a plant expressible promoter region;

(ii) said isolated DNA fragment comprising part of the parG encoding gene of said plant in direct orientation compared to the promoter region;

(iii) said isolated DNA fragment comprising part of the parG encoding gene of said plant in inverted orientation compared to the promoter region;

(iv) a 3' end region involved in transcription termination and polyadenylation;

(c) providing plant cells with said chimeric gene to create transgenic plant cells

(d) regenerating a population of transgenic plant lines from said transgenic plant cell; and

(e) identifying a stress tolerant plant line within said population of transgenic plant lines.

9. A DNA molecule comprising

- (i) a plant-expressible promoter
- (ii) a DNA region, which when transcribed yields an ParG inhibitory RNA molecule
- (iii) a 3' end region involved in transcription termination and polyadenylation, as described in any one of claims 1 to 8.

10. A plant cell comprising the DNA molecule of claims 9.

11. A plant consisting essentially of the plant cells of claim 10.

12. A process for producing stress tolerant plants, comprising the step of further crossing a plant of claim 11 with another plant.

13. Seeds and propagating material of a plant according to claim 10, comprising the chimeric gene of claim 9.

14. Plants obtained by the process of claim 8.

ABSTRACT

Methods and means are provided to increase the tolerance of plants to abiotic stress or adverse growing conditions, including drought, high light intensities, high temperatures, nutrient limitations and the like by reducing the activity of endogenous PARG proteins in plants.

1/3




Figure 1

2/3




Figure 2

3/3




Figure 3

SEQUENCE LISTING

<110> De Block, Marc

<120> Methods and means for increasing the tolerance of plants to stress conditions.

<130> BCS 03 2002

<160> 14

<170> PatentIn version 3.0

<210> 1

<211> 548

<212> PRT

<213> Arabidopsis thaliana

<400> 1

Met	Glu	Asn	Arg	Glu	Asp	Leu	Asn	Ser	Ile	Leu	Pro	Tyr	Leu	Pro	Leu
1				5					10					15	
Val	Ile	Arg	Ser	Ser	Ser	Leu	Tyr	Trp	Pro	Pro	Arg	Val	Val	Glu	Ala
			20					25					30		
Leu	Lys	Ala	Met	Ser	Glu	Gly	Pro	Ser	His	Ser	Gln	Val	Asp	Ser	Gly
		35					40					45			
Glu	Val	Leu	Arg	Gln	Ala	Ile	Phe	Asp	Met	Arg	Arg	Ser	Leu	Ser	Phe
	50					55					60				
Ser	Thr	Leu	Glu	Pro	Ser	Ala	Ser	Asn	Gly	Tyr	Ala	Phe	Leu	Phe	Asp
65					70					75					80
Glu	Leu	Ile	Asp	Glu	Lys	Glu	Ser	Lys	Arg	Trp	Phe	Asp	Glu	Ile	Ile
			85						90					95	
Pro	Ala	Leu	Ala	Ser	Leu	Leu	Leu	Gln	Phe	Pro	Ser	Leu	Leu	Glu	Val
			100					105						110	

His Phe Gln Asn Ala Asp Asn Ile Val Ser Gly Ile Lys Thr Gly Leu
 115 120 125
 Arg Leu Leu Asn Ser Gln Gln Ala Gly Ile Val Phe Leu Ser Gln Glu
 130 135 140
 Leu Ile Gly Ala Leu Leu Ala Cys Ser Phe Phe Cys Leu Phe Pro Asp
 145 150 155 160
 Asp Asn Arg Gly Ala Lys His Leu Pro Val Ile Asn Phe Asp His Leu
 165 170 175
 Phe Ala Ser Leu Tyr Ile Ser Tyr Ser Gln Ser Gln Glu Ser Lys Ile
 180 185 190
 Arg Cys Ile Met His Tyr Phe Glu Arg Phe Cys Ser Cys Val Pro Ile
 195 200 205
 Gly Ile Val Ser Phe Glu Arg Lys Ile Thr Ala Ala Pro Asp Ala Asp
 210 215 220
 Phe Trp Ser Lys Ser Asp Val Ser Leu Cys Ala Phe Lys Val His Ser
 225 230 235 240
 Phe Gly Leu Ile Glu Asp Gln Pro Asp Asn Ala Leu Glu Val Asp Phe
 245 250 255
 Ala Asn Lys Tyr Leu Gly Gly Gly Ser Leu Ser Arg Gly Cys Val Gln
 260 265 270
 Glu Glu Ile Arg Phe Met Ile Asn Pro Glu Leu Ile Ala Gly Met Leu
 275 280 285
 Phe Leu Pro Arg Met Asp Asp Asn Glu Ala Ile Glu Ile Val Gly Ala
 290 295 300
 Glu Arg Phe Ser Cys Tyr Thr Gly Tyr Ala Ser Ser Phe Arg Phe Ala
 305 310 315 320
 Gly Glu Tyr Ile Asp Lys Lys Ala Met Asp Pro Phe Lys Arg Arg Arg
 325 330 335
 Thr Arg Ile Val Ala Ile Asp Ala Leu Cys Thr Pro Lys Met Arg His
 340 345 350
 Phe Lys Asp Ile Cys Leu Leu Arg Glu Ile Asn Lys Ala Leu Cys Gly
 355 360 365
 Phe Leu Asn Cys Ser Lys Ala Trp Glu His Gln Asn Ile Phe Met Asp
 370 375 380
 Glu Gly Asp Asn Glu Ile Gln Leu Val Arg Asn Gly Arg Asp Ser Gly
 385 390 395 400
 Leu Leu Arg Thr Glu Thr Thr Ala Ser His Arg Thr Pro Leu Asn Asp
 405 410 415
 Val Glu Met Asn Arg Glu Lys Pro Ala Asn Asn Leu Ile Arg Asp Phe

420							425					430				
Tyr	Val	Glu	Gly	Val	Asp	Asn	Glu	Asp	His	Glu	Asp	Asp	Gly	Val	Ala	
		435					440					445				
Thr	Gly	Asn	Trp	Gly	Cys	Gly	Val	Phe	Gly	Gly	Asp	Pro	Glu	Leu	Lys	
	450					455					460					
Ala	Thr	Ile	Gln	Trp	Leu	Ala	Ala	Ser	Gln	Thr	Arg	Arg	Pro	Phe	Ile	
465					470					475					480	
Ser	Tyr	Tyr	Thr	Phe	Gly	Val	Glu	Ala	Leu	Arg	Asn	Leu	Asp	Gln	Val	
				485					490					495		
Thr	Lys	Trp	Ile	Leu	Ser	His	Lys	Trp	Thr	Val	Gly	Asp	Leu	Trp	Asn	
			500				505					510				
Met	Met	Leu	Glu	Tyr	Ser	Ala	Gln	Arg	Leu	Tyr	Lys	Gln	Thr	Ser	Val	
		515					520					525				
Gly	Phe	Phe	Ser	Trp	Leu	Leu	Pro	Ser	Leu	Ala	Thr	Thr	Asn	Lys	Ala	
	530					535					540					
Ile	Gln	Pro	Pro													
545																

<210> 2

<211> 169

<212> PRT

<213> Solanum tuberosum

<400> 2

Met 1	Glu	Asn	Arg	Glu 5	Asp	Val	Lys	Ser	Ile 10	Leu	Pro	Phe	Leu	Pro 15	Val
Cys	Leu	Arg	Ser 20	Ser	Ser	Leu	Phe	Trp 25	Pro	Pro	Leu	Val	Val 30	Glu	Ala
Leu	Lys	Ala 35	Leu	Ser	Glu	Gly	Pro 40	His	Tyr	Ser	Asn	Val 45	Asn	Ser	Gly
Gln 50	Val	Leu	Phe	Leu	Ala	Ile 55	Ser	Asp	Ile	Arg	Asn 60	Ser	Leu	Ser	Leu
Pro 65	Asp	Ser	Ser	Ile 70	Ser	Ser	Ser	Ala	Ser	Asp 75	Gly	Phe	Ser	Leu 80	Leu
Phe	Asp	Asp	Leu	Ile 85	Pro	Arg	Asp	Glu	Ala 90	Val	Lys	Trp	Phe	Lys 95	Glu
Val	Val	Pro	Lys 100	Met	Ala	Asp	Leu	Leu 105	Leu	Arg	Leu	Pro	Ser 110	Leu	Leu

Glu Ala His Tyr Glu Lys Ala Asp Gly Gly Ile Val Lys Gly Val Asn
 115 120 125
 Thr Gly Leu Arg Leu Leu Glu Ser Gln Gln Pro Gly Ile Val Phe Leu
 130 135 140
 Ser Gln Glu Leu Val Gly Ala Leu Leu Ala Cys Ser Phe Phe Cys Tyr
 145 150 155 160
 Ser Leu Pro Met Ile Glu Val Ser Val
 165

<210> 3

<211> 1647

<212> DNA

<213> Arabidopsis thaliana

<400> 3
 atggagaatc gcgaagatct taactcaatt ctcccgtaac ttccacttgt aatcggttcg 60
 tcgtcgctgt attggccgcc gcgtgtggtg gaggcgttaa aggcaatgtc tgaaggacca 120
 tctcacagcc aagttgactc aggagaggtt ctacggcaag ctattttcga tatgagacga 180
 tccttatctt tctctactct cgagccatct gcttctaatt gctacgcatt tctctttgac 240
 gaattgattg atgagaaaga atcaaagaga tggttcgatg agattatccc agcattggcg 300
 agcttacttc tacagtttcc atctctgtta gaagtgcatt tccaaaatgc tgataatatt 360
 gttagtggaa tcaaaaccgg tcttcgtttg ttaaattccc aacaagctgg cattgttttc 420
 ctcagccagg agttgattgg agctcttctt gcatgctctt tcttttgttt gtttcgggat 480
 gataatagag gtgcaaaaca ccttcagtc atcaactttg atcatttggt tgcaagcctt 540
 tatataagtt atagtcaaag tcaagaaagc aagataagat gtattatgca ttactttgaa 600
 aggttttgct cctgcgtgcc tattggtatt gtttcatttg aacgcaagat taccgctgct 660
 cctgatgctg atttctggag caagtctgac gtttctcttt gtgcatttaa ggttcaactt 720
 tttggggttaa ttgaagatca acctgacaat gctctcgaag tggactttgc aaacaagtat 780
 ctccggaggtg gttccctaag tagagggtgc gtgcaggaag agatacgctt catgattaac 840
 cctgaattaa tcgctggcat gcttttcttg cctcggatgg atgacaatga agctatagaa 900
 atagttggtg cggaaagatt ttcattgtac acagggtatg catcttcggt tcggtttgct 960
 ggtgagtaca ttgacaaaaa ggcaatggat cctttcaaaa ggccaagaac cagaattggt 1020

gcaattgatg cattatgtac accgaagatg agacacttta aagatatatg tcttttaagg 1080
gaaattaata aggcactatg tggcttttta aattgtagca aggccttggga gcaccagaat 1140
atattcatgg atgaaggaga taatgaaatt cagcttgtcc gaaacggcag agattctggg 1200
cttctgcgta cagaaactac tgcgtcacac cgaactccac taaatgatgt tgagatgaat 1260
agagaaaagc ctgctaacaa tcttatcaga gatttttatg tggaggagt tgataacgag 1320
gatcatgaag atgatgggtg cgcgacaggg aattggggat gtggtgtttt tggaggagac 1380
ccagagctaa aggctacgat acaatggctt gctgcttccc agactcgaag accatttata 1440
tcatattaca cctttggagt agaggcactc cgaaacctag atcaggtgac gaagtggatt 1500
ctttcccata aatggactgt tggagatctg tggacatga tgtagaata ttctgtctaa 1560
aggctctaca agcaaaccag tgttggtctc tttctctggc tacttccatc tctagctacc 1620
accaacaaag ctatccagcc gccttga 1647

<210> 4

<211> 598

<212> DNA

<213> Solanum tuberosum

<400> 4

gcaatggaga atagagaaga cgtgaagtca atccttccct ttttgccggg gtgtctccga 60
tcatcttctc ttttctggcc gccgctagtt gttgaagcac tgaaagccct ctctgaaggc 120
cctcattaca gcaatgttaa ctccggccaa gtctcttcc tcgcaatctc cgacattcgg 180
aattcccttt cactacctga ttcttcaatt tctctctctg cttcagacgg attttctctc 240
ttatttgatg atttaattcc tagggatgaa gctgttaaag ggttcaaaga agtgggtgccg 300
aaaatggcgg atttgctatt gcggttgccct tccttattgg aggcacta tgagaaggct 360
gatgggtggaa ttgttaaagg agtcaacact ggtcttctgct tattggaatc acaacagcct 420
ggcattgttt tcttcagtcg ggaattagtc ggtgctcttc ttgcatgttc cttcttttgc 480
tattccctac caatgataga ggtatctgta tgatcagtat gacgagaaat ttgaaaataa 540
attgaagtgc attcttcact attttgagag gattggctca ttgatactg cgggctac 598

<210> 5

<211> 37

<212> DNA

<213> Artificial

<220>

<223> oligonucleotide primer ParGAt1

<400> 5
ggatccctg caggacaaaa aggcaatgga tcctttc

37

<210> 6

<211> 39

<212> DNA

<213> Artificial

<220>

<223> oligonucleotide primer ParGAt2

<400> 6
gcacgaattc gcggcgcggtg tgctcccaag ccttgctac

39

<210> 7

<211> 39

<212> DNA

<213> Artificial

<220>

<223> oligonucleotide primer ParGSt1

<400> 7
ggatccctg caggctcact atgagaaggc tgatggtgg

39

<210> 8

<211> 43

<212> DNA

<213> Artificial

<220>

<223> oligonucleotide primer ParGSt2

<400> 8

gcacgaattc gcggccgcgt catactgatac atacagatac etc

43

<210> 9

<211> 13466

<212> DNA

<213> Artificial

<220>

<223> nucleotide sequence of pTVB428

<220>

<221> misc_feature

<222> (198)..(222)

<223> Right T-DNA border

<220>

<221> misc_feature

<222> (983)..(273)

<223> 3' ocs (3' untranslated end of octopine synthase gene)

<220>

<221> misc_feature

<222> (995)..(1155)

<223> part of poly (ADP-ribose) glycohydrolase

<220>

<221> misc_feature

<222> (1929)..(1188)

<223> intron 2 from the Pdk gene of Flaveria

<220>

<221> misc_feature

<222> (2122)..(1962)

<223> part of poly (ADP-ribose) glycohydrolase

<220>

<221> misc_feature

<222> (3476)..(2131)

<223> 35S promoter region from Cauliflower Mosaic Virus

<220>

<221> misc_feature

<222> (3948)..(3737)

<223> 3' untranslated end of gene 7 from Agrobacterium tumefaciens

<220>

<221> misc_feature

<222> (4521)..(3970)

<223> bar coding region

<220>

<221> misc_feature

<222> (6247)..(4522)

<223> PSSuAra promoter region

<220>

<221> misc_feature

<222> (6415)..(6439)

<223> left border region of T-DNA of Agrobacterium tumefaciens

```

<400> 9
agattcgaag ctcggtcccg tgggtgttct gtcgtctcgt tgtacaacga aatccattcc      60
cattccgcgc tcaagatggc ttcccctcgg cagttcatca gggctaaatc aatctagccg      120
acttgtccgg tgaatgggc tgcactccaa cagaaacaat caaacaacaa tacacagcga      180
cttattcaca cgcgacaaat tacaacggta tatatcctgc cagtactcgg ccgtcgaccg      240
cgggtaccccg gaattaagct tgcattgctg caggtcctgc tgagcctcga catgttgtcg      300
caaaattcgc cctggaccgg cccaacgatt tgtcgtcact gtcaagggtt gacctgcact      360
tcatttgggg ccacataca ccaaaaaaat gctgcataat tctcggggca gcaagtcggg      420
taccggggcg ccgtgctgga ccgggttgaa tgggtgcccg aactttcggg agagcggacg      480
gccaatactc aatttcaagg aatctcaccg atgcgcgccg gcgggggaacc ggagttccct      540
tcagtgaacg ttattagttc gccgctcggg gtgtcgtaga tactagcccc tggggccttt      600
tgaaatttga ataagattta tgtaatcagt cttttaggtt tgaccgggtt tgccgctttt      660
tttaaaattg gatttgtaat aataaaacgc aattgtttgt tattgtggcg ctctatcata      720
gatgtcgcta taaacctatt cagcacataa tattgttttc attttaatat tgtacatata      780
agtagtaggg tacaatcagt aaattgaacg gagaatatta ttcataaaaa tacgatagta      840
acgggtgata tattcattag aatgaaccga aaccggcggg aaggatctga gctacacatg      900
ctcagggttt ttacaacgtg cacaacagaa ttgaaagcaa atatcatgcg atcataggcg      960
tctcgcatat ctcattaaag caggactcta gagacaaaaa ggcaatggat cctttcaaaa     1020
ggcgaagaac cagaattggt gcaattgatg cattatgtac accgaagatg agacacttta     1080
aagatatatg tcttttaagg gaaattaata aggcactatg tggcttttta aattgtagca     1140
aggcttggga gcaccatcga ttctgaaccc agcttcccaa ctgtaatcaa tccaaatgta     1200
agatcaatga taacacaatg acatgatcta tcatgttacc ttgtttattc atgttcgact     1260
aattcattta attaatagtc aatccattta gaagttaata aaactacaag tattatttag     1320
aaattaataa gaatgttgat tgaaaataat actatataaa atgatagatc ttgcgctttg     1380
ttatattagc attagattat gttttgttac attagattac tgtttctatt agtttgatat     1440

```

tatttggttac tttagcttgt tatttaatat tttggtttatt gataaattac aagcagattg 1500
gaattttctaa caaaatattt attaacctttt aaactaaaat atttagtaat ggtatagata 1560
tttaattata taataaacta ttaatcataa aaaaatatta ttttaattta tttattotta 1620
tttttactat agtattttat cattgatatt taattcatca aaccagctag aattactatt 1680
atgattaataa caaatattaa tgctagtata tcatcttaca tgttcgatca aattcattaa 1740
aaataatata cttactctca actttttatct tcttcgtctt acacatcaact tgtcatattt 1800
ttttacatta ctatgttggt tatgtaaaca atatatttat aaattatttt ttcacaatta 1860
taacaactat attattataa tcataactaat taacatcaact taactatttt atactaaaag 1920
gaaaaaagaa aataattatt tccctaccaa gctgggggtac cgggtgctccc aagccttgct 1980
acaatttaaa aagccacata gtgccttatt aatttcctt aaaagacata tatcttttaa 2040
gtgtctcatc ttgggtgtac ataatgcac aattgcaaca attctgggtc ttgcctttt 2100
gaaaggatcc attgcctttt tgtcctcgag cgtgtcctct ccaaagaaa tgaacttcct 2160
tatatagagg aagggtcttg cgaaggatag tgggattgtg cgtcatccct tacgtcagt 2220
gagatgtcac atcaatccac ttgctttgaa gacgtgggtg gaacgtcttc ttttccacg 2280
atgctcctcg tgggtggggg tccatctttg ggaccactgt cggcagagag atcttgaatg 2340
atagcctttc ctttatcgca atgatggcat ttgtaggagc caccttcctt ttctactgtc 2400
ctttcgatga agtgacagat agctgggcaa tggaaatccga ggaggtttcc cgaaattatc 2460
ctttgttgaa aagtctcaat agccctttgg tcttctgaga ctgtatcttt gacatttttg 2520
gagtagacca gagtgtcgtg ctccaccatg ttgaagaaga ttttcttctt gtcattgagt 2580
cgtaaaagac tctgtatgaa ctgttcgcca gtcttcacgg cgagtctctg tagatcctcg 2640
atttgaatct tagactccat gcatggcctt agattcagta ggaactacct ttttagagac 2700
tccaatctct attacttgcc ttgggttatg aagcaagcct tgaatcgtcc atactggaat 2760
agtacttctg atcttgagaa atatgtcttt ctctgtgttc ttgatgcaat tagtcctgaa 2820
tcttttgact gcatctttaa ccttcttggg aagggtatttg atctcctgga gattgttact 2880
cgggtagatc gtcttgatga gacctgotgc gtaggcctct ctaaccatct gtgggtcagc 2940
attctttctg aaattgaaga ggctaacctt ctcatatca gtgggtgaaca tagtgtcgtc 3000
accttcacct tcgaacttcc ttcttagatc gtaaagatag aggaaatcgt ccattgtaat 3060
ctccggggca aaggagatct cttttggggc tggatcactg ctgggccttt tggttcctag 3120
cgtgagccag tgggcttttt gctttgggtg gcttggtagg gccttagcaa agctcttggg 3180

cttgagttga gcttctcctt tggggatgaa gttcaacctg tctgtttgct gacttgttgt 3240
gtacgcgtca gctgctgctc ttgcctctgt aatagtggca aatttcttgt gtgcaactcc 3300
gggaacgccg tttgttgccg cctttgtaca accccagtca tcgtatatac cggcatgtgg 3360
accgttatat acaacgtagt agttgatatg aggggtgtga ataccgatt ctgctctgag 3420
aggagcaact gtgctgttaa gctcagattt ttgtgggatt ggaattaatt cgtcgagcgg 3480
ccgctcgacg agcgcgcga tatcgcgatc gcccgggccg gccatttaaa tgaattcgag 3540
ctcgggtacc aaacgcggcc gcaagctata acttcgtata gcatacatta tacgaagtta 3600
ttcgactcta gaggatccca attcccatgc atggagtcaa agattcaa at agaggacact 3660
tctcgaactc ggccgtcgaa ctcggccgtc gagtacatgg tcgataagaa aaggcaattt 3720
gtagatgtta attcccatct tgaaagaa atagtttaaa tatttattga taaaataaca 3780
agtcaggat tatagtccaa gcaaaaacat aaatttattg atgcaagttt aaattcagaa 3840
atatttcaat aactgattat atcagctggc acattgccgt agatgaaaga ctgagtgcga 3900
tattatgtgt aatacataaa ttgatgatat agctagctta gctcatcggg ggatcctaga 3960
cgctgagat cagatctcgg tgacgggcag gaccggacgg ggccgtaccg gcaggctgaa 4020
gtccagctgc cagaaaccca cgtcatgcc a gttccgtgc ttgaagccgg ccgcccgcag 4080
catgccgcgg ggggcatatc cgagcgcctc gtgcatgcgc acgctcgggt cgttgggcag 4140
cccgatgaca gcgaccacgc tcttgaagcc ctgtgcctcc agggacttca gcagggtgggt 4200
gtagagcgtg gagccagtc ccgtccgctg gtggcggggg gagacgtaca cggtcgactc 4260
ggccgtccag tcgtaggcgt tgcgtgcctt ccaggggccc gcgtaggcga tgccggcgac 4320
ctcgccgtcc acctcggcga cgagccaggg atagcgctcc cgcagacgga cgaggctcgtc 4380
cgctccactc tcgggttcct gcggctcgggt acggaagttg accgtgcttg tctcgatgta 4440
gtgggtgacg atgggtgcga ccgcggcat gtccgcctcg gtggcacggc ggatgtcggc 4500
cgggcgtcgt tctgggtcca ttgttcttct ttactctttg tgtgactgag gtttgggtcta 4560
gtgcttttgg catctatata taatgataac aacaatgaga acaagctttg gagtgatcgg 4620
agggcttagg atacatgaga ttcaagtgga ctaggatcta caccgttggg ttttgagtgt 4680
ggatatgtgt gaggttaatt ttacttggta acggccacaa aggcctaagg agagggtgtt 4740
agacccttat cggcttgaac cgctggaata atgccacgtg gaagataatt ccatgaatct 4800
tatcgttatc tatgagtga attgtgtgat ggtggagtgg tgcttgctca ttttacttgc 4860
ctgggtggact tggcccttct cttatgggga atttatattt tacttactat agagctttca 4920

tacctttttt ttaccttgga tttagttaat atataatggt atgattcatg aataaaaaatg 4980
ggaaattttt gaatttgtac tgctaaatgc ataagattag gtgaaactgt ggaatatata 5040
tttttttcat ttaaaagcaa aatttgcctt ttactagaat tataaatata gaaaaatata 5100
taacattcaa ataaaaatga aaataagaac tttcaaaaaa cagaactatg tttaatgtgt 5160
aaagattagt cgcacatcaa gtcactgtt acaatatgtt acaacaagtc ataagcccaa 5220
caaagttagc acgtctaaat aaactaaaga gtccacgaaa atattacaaa tcataagccc 5280
aaciaaagtta ttgatcaaaa aaaaaaacg cccaacaaag ctaaacaaag tccaaaaaaa 5340
acttctcaag tctccatctt cctttatgaa cattgaaaac tatacacaaa acaagtcaga 5400
taaactctctt tctgggcctg tcttcccaac ctctacatc acttccctat cggattgaat 5460
gttttacttg taccttttcc gttgcaatga tattgatagt atgtttgtga aaactaatag 5520
ggttaacaat cgaagtcag gaatatggat ttgggtccaag attttccgag agctttctag 5580
tagaaagccc atcaccagaa atttactagt aaaataaatc accaattagg tttcttatta 5640
tgtgccaaat tcaatataat tatagaggat atttcaaag aaaacgtatg aatgttatta 5700
gtaaatggtc aggtaaagaca ttaaaaaaat cctacgtcag atattcaact ttaaaaaattc 5760
gatcagtggt gaattgtaca aaaatttggg atctactata tatatataat gctttacaac 5820
acttggattt ttttttggag gctggaattt ttaatctaca tatttgtttt ggccatgcac 5880
caactcattg tttagtgtaa tactttgatt ttgtcaaata tatgtgttcg tgtatatttg 5940
tataagaatt totttgacca tatacacaca cacatatata tatatatata tatattatat 6000
atcatgcact ttaattgaa aaaataatat atatatatat agtgcatttt ttctaacaac 6060
catatatggt gcgattgatc tgcaaaaata ctgctagagt aatgaaaaat ataacttatt 6120
gctgaaatta tctcagatgt taagattttt ttaaagtaaa ttctttcaaa ttttagctaa 6180
aagtcttgta ataactaaag aataatacac aatctcgacc acggaaaaaa aacacataat 6240
aaatttgaat ttcgaccgag gtacccggaa ttgggttata attacctcag gtcgaggaat 6300
taattcggtta cgtacctaat aacttcgtat agcatacatt atacgaagtt atatggatct 6360
cgaggcatta cggcattacg gcaactcgga ggggtcccaat tcgagcatgg agccatttac 6420
aattgaatat atcotgccc cgctgccgct ttgcaccggg tggagcttgc atgttggttt 6480
ctacgcagaa ctgagccggt taggcagata atttccattg agaactgagc catgtgcacc 6540
ttccccccaa cacgggtgagc gacggggcaa cggagtgatc cacatgggac ttttaaacat 6600
catccgtcgg atggcggtgc gagagaagca gtcgatccgt gagatcagcc gacgcaccgg 6660

gcaggcgcgc aacacgatcg caaagtatct gaacgcaggt acaatcgagc cgacgttcac 6720
ggtagcggaa cgaccaagca agctagctta gtaaagccct cgctagatct taatgcggat 6780
gttgcgatta ctctcgcaac tattgcgata acaagaaaaa gccagcctct catgatatat 6840
ctcccaatct gtgtagggct tattatgcac gcttaaaaaat aataaaagca gacttgacct 6900
gatagtttgg ctgtgagcaa ttatgtgctt agtgcattct acgcttgagt taagccgcgc 6960
cgcaagcgg cgctcgcttg aacgaattgt tagacattat ttgccgacta ccttggtgat 7020
ctcgcccttc acgtagtggg caaattcttc caactgatct gcgcgcgagg ccaagcgatc 7080
ttcttcttgt ccaagataag cctgtctagc ttcaagtatg acgggctgat actgggcccgg 7140
caggcgctcc attgccagct cggcagcgac atccttcggc gcgattttgc cggttactgc 7200
gctgtaccaa atgcgggaca acgtaagcac tacatttcgc tcatcgccag ccagtcggg 7260
cggcgagttc catagcgtta aggtttcatt tagcgcctca aatagatcct gttcaggaac 7320
cggatcaaag agttcctccg ccgctggacc taccaaggca acgctatggt ctcttgcttt 7380
tgtcagcaag atagccagat caatgtcgat cgtggctggc tcgaagatac ctgcaagaat 7440
gtcattgcgc tgccattctc caaattgcag ttcgcgctta gctggataac gccacggaat 7500
gatgtcgtcg tgcacaacaa tgggtgaactc tacagcgcgg agaattctgc tctctccagg 7560
ggaagccgaa gtttccaaaa ggtcggtgat caaagctcgc cgcgttggtt catcaagcct 7620
tacggtcacc gtaaccagca aatcaatctc actgtgtggc ttcaggccgc catccactgc 7680
ggagccgtac aaatgtacgg ccagcaacgt cggttcgaga tggcgctcga tgacgccaac 7740
tacctctgat agttgagtcg atacttcggc gatcaccgct tccctcatga tgtttaactt 7800
tggttttaggg cgactgcctt gctgcgtaac atcgttgctg ctccataaca tcaaaccatcg 7860
acccacggcg taacgcgctt gctgcttggg tgcccagagg atagactgta ccccaaaaaa 7920
acagtcataa caagccatga aaaccgccac tgcgcgctta ccaccgctgc gttcgggtcaa 7980
ggttctggac cagttgcgtg agcgcatacg ctacttgcat tacagcttac gaaccgaaca 8040
ggcttatgtc cactgggttc gtgccttcat ccgtttccac ggtgtgcgtc acccggaac 8100
cttgggcagc agcgaagtcg aggcatttct gtccctggctg gcgaacgagc gcaaggtttc 8160
ggctctccacg catcgtcagg cattggcggc cttgctgttc ttctacggca agtgctgtgc 8220
acggatctgc cctggcttca ggagatcgga agacctcggc cgtccgggcg cttgcccgtg 8280
gtgctgacc cggatgaagt ctctagagct ctagagggtt cgcattcctcg gttttctgga 8340
aggcgagcat cgtttgttcg ccagcttct gtatggaacg ggcattcgga tcagtgggg 8400

tttgcaactg cgggtcaagg atctggattt cgatcacggc acgatcatcg tgcgggaggg 8460
caagggctcc aaggatcggg ccttgatgtt acccgagagc ttggcaccca gcctgcgcga 8520
gcagggatcg atccaacccc tcegtgcta tagtgacgtc ggcttctgac gttcagtgca 8580
gocgtcttct gaaaacgaca tgcgcacaa gtcctaagtt acgcgacagg ctgcgcct 8640
gcccttttcc tggcgtttcc ttgtcgcgtg ttttagtcgc ataaagtaga atacttgca 8700
ctagaaccgg agacattacg ccatgaacaa gagcgccgcc gctggcctgc tgggctatgc 8760
ccgcgtcagc accgacgacc aggacttgac caaccaacgg gccgaactgc acgcggccgg 8820
ctgcaccaag ctgttttccg agaagatcac cggcaccagg cgcgaccgcc cggagctggc 8880
caggatgctt gaccacctac gccctggcga cgttgtgaca gtgaccaggc tagaccgcct 8940
ggcccgacgc acccgcgacc tactggacat tgccgagcgc atccaggagg ccggcgccggg 9000
cctgcgtagc ctggcgagac cgtgggcccga caccaccaac ccggccggcc gcattggtgtt 9060
gaccgtgttc gcgggcattg ccgagttcga gcgttcccta atcatcgacc gcacccggag 9120
cgggcgcgag gccgccaagg ccgaggcgt gaagtttggc cccgccccta cctcaccacc 9180
ggcacagatc gcgcacgccc gcgagctgat cgaccaggaa ggccgcaccg tgaaagaggc 9240
ggctgcactg cttggcgtgc atcgctcgac cctgtaccgc gcacttgagc gcagcgagga 9300
agtgcgccc accgaggcca ggccggcgcg tgccttccgt gaggacgcatt tgaccgaggc 9360
cgacgccttg gcggccgccc agaataaacg ccaagaggaa caagcatgaa accgcaccag 9420
gacggccagg acgaaccgtt tttcattacc gaagagatcg aggcggagat gatcgcgccc 9480
gggtacgtgt tcgagccgcc cgcgcacgtc tcaaccgtgc ggctgcatga aatcctggcc 9540
ggtttgtctg atgccaaagt ggccggcctgg ccggccagct tggccgctga agaaaccgag 9600
cgccgcctgc taaaaaggtg atgtgtattt gagtaaaaca gcttgctca tgcggtcgt 9660
gcgtatatga tgcgatgagt aaataaaca atacgcaagg ggaacgcatt aaggttatcg 9720
ctgtacttaa ccagaaaggc gggtcaggca agacgaccat cgcaacccat ctgcccgcg 9780
ccctgcaact cgcggggggc gatgttctgt tagtcgattc cgatccccag ggcagtgcc 9840
gcgattgggc ggccgtgcg gaagatcaac cgctaaccgt tgcggcacc gaccgcccga 9900
cgattgaccg cgacgtgaag gccatcgccc ggccgcactt cgtagtgate gacggagcgc 9960
cccaggcggc ggacttggt gtgtccgcga tcaaggcagc cgacttcgtg ctgattccgg 10020
tgcagccaag cccttacgac atatgggcca ccgccgacct ggtggagctg gtttaagcagc 10080
gcattgaggt cacggatgga aggctacaag cggcctttgt cgtgtcgcg gcgatcaaag 10140

gcacgcgcacat cggcggtgag gttgccgagg cgctggccgg gtacgagctg cccattcttg 10200
agtecccgatat cacgcagcgc gtgagctacc caggcactgc cgccgccggc acaaccgttc 10260
ttgaatcaga acccgagggc gacgctgccc gcgaggtcca ggcgctggcc gctgaaatta 10320
aatcaaaact catttgagtt aatgaggtaa agagaaaatg agcaaaagca caaacacgct 10380
aagtgccggc cgtccgagcg cacgcagcag caaggctgca acgttgGCCa gcctggcaga 10440
cacgccagcc atgaagcggg tcaactttca gttgccggcg gaggatcaca ccaagctgaa 10500
gatgtacgcg gtacgccaaG gcaagaccat taccgagctg ctatctgaat acatcgcgca 10560
gctaccagag taaatgagca aatgaataaa tgagtagatg aatttttagcg gctaaaggag 10620
gcggcatgga aaatcaagaa caaccaggca ccgacgccgt ggaatgcccc atgtgtggag 10680
gaaeggcgcg ttggccaggc gtaagcggct gggttgtctg ccggccctgc aatggcactg 10740
gaacccccaG gcccgaggaa tcggcgtagc ggtcgcaaac catccggccc ggtacaaatc 10800
ggcgcgggcg tgggtgatga cotgggtggag aagttgaagg ccgcgagggc cgcccagcgg 10860
caacgcacgc aggcagaagc acgccccggt gaatcgtagc aagcggccgc tgatcgaatc 10920
cgcaaagaat cccggcaacc gccggcagcc ggtgcgccgt cgattaggaa gccgccaag 10980
ggcgacgagc aaccagattt tttcggtccg atgctctatg acgtgggcac ccgcgatagt 11040
cgcagcatca tggacgtggc cgttttccgt ctgtcgaagc gtgaccgacg agctggcgag 11100
gtgatccgct acgagcttcc agacgggcac gtagaggttt ccgcagggcc ggccggcatg 11160
gccagtgtgt gggattacga cctggtagtG atggcggttt cccatctaac cgaatccatg 11220
aaccgatacc gggaagggaa gggagacaag cccggccgcg tgttccgtcc acacgttgcg 11280
gacgtactca agttctgccg gcgagccgat ggcggaaagc agaaagacga cctggtagaa 11340
acctgcattc ggttaaacac cacgcacgtt gccatgcagc gtacgaagaa ggccaagaac 11400
ggccgcctgg tgacggtatc cgagggtgaa gccttgatta gccgctacaa gatcgtaaag 11460
agcgaaccg ggccggccga gtacatcgag atcgagctag ctgattggat gtaccgcgag 11520
atcacagaag gcaagaaccc ggacgtgctg acggttcacc ccgattactt tttgatcgat 11580
cccggcatcg gccgttttct ctaccgcctg gcacgcgcg ccgcaggcaa ggcagaagcc 11640
agatggttgt tcaagacgat ctacgaacgc agtggcagcg ccggagagtt caagaagttc 11700
tgtttcaccg tgcgcaagct gatcgggtca aatgacctgc cggagtacga tttgaaggag 11760
gaggcggggc aggctggccc gatcctagtc atgcgtacc gcaacctgat cgagggcgaa 11820
gcatccgcg gttcctaattg tacggagcag atgctagggc aaattgccct agcaggggaa 11880

aaaggtcgaa aaggtctctt tctgtggat agcacgtaca ttgggaaccc aaagccgtac 11940
attgggaacc ggaaccgta cattgggaac ccaaagccgt acattgggaa ccggtcacac 12000
atgtaagtga ctgatataaa agagaaaaaa ggcgattttt ccgcctaaaa ctctttaaaa 12060
cttattaaaa ctcttaaaac ccgcctggcc tgtgcataac tgtctggcca gcgcacagcc 12120
gaagagctgc aaaaagcgcc tacccttcgg tgcgtgcgt ccctacgccc cgccgcttcg 12180
cgteggccta tcgcgccgc tggccgctca aaaatggctg gcctacggcc aggcaatcta 12240
ccagggcgcg gacaagccgc gccgtcgcca ctgcaccgcc ggcgcccaca tcaaggcacc 12300
ctgcctcgcg cgtttcggtg atgacggtga aaacctctga cacatgcagc tcccgagac 12360
ggtcacagct tgtctgtaag cggatgccgg gagcagacaa gcccgtcagg gcgcgtcagc 12420
gggtgttggc gggtgtcggg gcgcagccat gaccagtcga cgtagcgata gcggagtgt 12480
tactggctta actatgcggc atcagagcag attgtactga gagtgcacca tatgcggtgt 12540
gaaataccgc acagatgcgt aaggagaaaa taccgcatca ggcgctcttc cgcttctctg 12600
ctcactgact cgctgcgctc ggtcgttcgg ctgcggcgag cggatatcagc tcaactcaaag 12660
gcggtaatac gggtatccac agaatcaggg gataacgcag gaaagaacat gtgagcaaaa 12720
ggccagcaaa aggccaggaa ccgtaaaaag gccgcgttgc tggcggtttt ccataggctc 12780
cgccccctg acgagcatca caaaaatcga cgctcaagtc agaggtggcg aaacccgaca 12840
ggactataaa gataccaggc gtttccccct ggaagctccc tcgtgcgctc tctgttccg 12900
accctgcgc ttaccggata cctgtccgc tttctccctt cgggaagcgt ggcgctttct 12960
catagctcac gctgtaggta tctcagttcg gtgtaggctg ttcgctccaa gctgggctgt 13020
gtgcacgaac ccccggttca gcccgaccgc tgcgccttat ccggttaacta tcgtcttgag 13080
tccaacccgg taagacacga cttatcgcca ctggcagcag cactggtaa caggattagc 13140
agagcgaggt atgtaggcgg tgctacagag ttcttgaagt ggtggcctaa ctacggctac 13200
actagaagga cagtatttgg tatctgcgct ctgctgaagc cagttacctt cggaaaaaga 13260
gttggtagct cttgatccgg caaacaacc accgctggta gcggtggtt tttgtttgc 13320
aagcagcaga ttacgcgcag aaaaaaagga tctcaagaag atccggaaaa cgcaagcgca 13380
aagagaaagc aggtagcttg cagtgggctt acatggcgat agctagactg ggcggtttta 13440
tggacagcaa gcgaaccgga attgcc 13466

<210> 10

<211> 31

<212> PRT

<213> Artificial

<220>

<223> Consensus sequence 1 of PARG protein

<220>

<221> VARIANT

<222> (2)..(2)

<223> X represents any amino acid

<220>

<221> VARIANT

<222> (8)..(10)

<223> X represents any amino acid

<220>

<221> VARIANT

<222> (14)..(17)

<223> X represents any amino acid

<220>

<221> VARIANT

<222> (19)..(19)

<223> X represents any amino acid

<220>

<221> VARIANT

<222> (27)..(29)

<223> X represents any amino acid

<400> 10

Leu Xaa Val Asp Phe Ala Asn Xaa Xaa Xaa Gly Gly Gly Xaa Xaa Xaa
1 5 10 15

Xaa Gly Xaa Val Gln Glu Glu Ile Arg Phe Xaa Xaa Xaa Pro Glu
20 25 30

<210> 11

<211> 20

<212> PRT

<213> Artificial

<220>

<223> Consensus sequence 2 for PARG protein

<220>

<221> VARIANT

<222> (3)..(3)

<223> X represents any amino acid

<220>

<221> VARIANT

<222> (10)..(10)

<223> X represents any amino acid

<220>

<221> VARIANT

<222> (13)..(14)

<223> X represents any amino acid

<220>

<221> VARIANT

<222> (17)..(19)

<223> X represents any amino acid

<400> 11

Thr Gly Xaa Trp Gly Cys Gly Ala Phe Xaa Gly Asp Xaa Xaa Leu Lys
1 5 10 15

Xaa Xaa Xaa Gln
20

<210> 12

<211> 13

<212> PRT

<213> Artificial

<220>

<223> Consensus sequence 3 for PARG protein

<220>

<221> VARIANT

<222> (2)..(4)

<223> X represents any amino acid

<220>

<221> VARIANT

<222> (6)..(9)

<223> X represents any amino acid

<400> 12

Asp Xaa Xaa Xaa Arg Xaa Xaa Xaa Xaa Ala Ile Asp Ala
1 5 10

<210> 13

<211> 10

<212> PRT

<213> Artificial

<220>

<223> Consensus sequence 4 for PARG protein

<220>

<221> VARIANT

<222> (3) .. (4)

<223> X represents any amino acid

<220>

<221> VARIANT

<222> (7) .. (8)

<223> X represents any amino acid

<400> 13

Arg	Glu	Xaa	Xaa	Lys	Ala	Xaa	Xaa	Gly	Phe
1				5					10

<210> 14

<211> 11

<212> PRT

<213> Artificial

<220>

<223> conserved PARG region

<220>

<221> VARIANT

<222> (2) .. (5)

<223> X represents any amino acid

<220>

<221> VARIANT

<222> (2)..(5)

<223> X represents any amino acid

<400> 14

Gly	Xaa	Xaa	Xaa	Xaa	Ser	Xaa	Tyr	Thr	Gly	Tyr
1				5					10	

PO#EP004003995



This Page is inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURED OR ILLEGIBLE TEXT OR DRAWING
- ☒ SKEWED/SLANTED IMAGES
- ☐ COLORED OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REPERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images
problems checked, please do not report the
problems to the IFW Image Problem Mailbox**